

Fuzzy Dark Matter at neutrino experiments

based on :

VB, J. Kopp, J. Liu, P. Prass, X. Wang arXiv:1705.09455


Vedran Brdar




Johannes Gutenberg Universität Mainz

NUFACT 2017

Dark Matter - the mystery of this century




- ▶ not charged under $U(1)_{EM}$ and $SU(3)_C$
- ▶ stable or long lived
- ▶ not in SM particle list



- ▶ Direct detection
 - ▶ nuclear recoils from DM scattering
- ▶ Collider searches
 - ▶ typical signal: missing energy + mono object
- ▶ Indirect detection
 - ▶ classified by annihilation product: $\gamma, \nu, e^+ \dots$

Dark Matter - vast number of candidates



- ▶ in this talk we will focus on the low end of DM spectrum – fuzzy DM with mass $\mathcal{O}(10^{-22})$ eV
- ▶ we consider both scalar and vector ultralight DM

Properties of Fuzzy DM candidate

Fuzzy DM can address:

- ▶ “core vs. cusp problem” – DM density profile discrepancy between measurements and simulations

→ DM delocalization

(huge Compton wave length $\lambda = 2\pi/m_\phi \simeq 0.4 \text{ pc} \times (10^{-22} \text{ eV}/m_\phi)$)

- ▶ “missing satellites problem” – lower than expected abundance of dwarf galaxies

→ higher probability for tidal disruption of DM subhalos and suppression of the matter power spectrum at small scales (Hui et al. 1610.08297)

- ▶ “too big to fail problem” – apparent failure of many of the most massive Milky Way subhalos to host visible dwarf galaxies

→ Fuzzy DM predicts fewer such subhalos (Marsh et al. 1307.1705)

- ▶ admittedly, better treatment of baryonic physics in simulations (1602.05957, 1202.0554) may solve these puzzles but the possibility that DM physics plays a crucial role is not excluded

DM production

Misalignment mechanism

Arias et al. 1201.5902
Nelson & Scholtz 1105.2812
Golovnev et al. 0802.2068

- ▶ EOM for real scalar field ϕ

$$\ddot{\phi} + 3H\dot{\phi} + m_\phi^2\phi = 0.$$

- ▶ while $3H \gg m_\phi$, ϕ is “frozen”
- ▶ at $3H = m_\phi$ damping term stops dominating and the field can start to oscillate
- ▶ for vector DM ϕ^μ one introduces coupling to gravity $\sim R\phi_\mu\phi^\mu$
- ▶ The mass of ϕ^μ can be generated either through the Stückelberg mechanism or from spontaneous symmetry breaking in a dark Higgs sector
- ▶ we consider both polarized and unpolarized vector DM (polarization may be altered during structure formation)

Model

Relevant part of the Lagrangian:

$$\text{Scalar } \mathcal{L}_{\text{scalar}} = \bar{\nu}_L^\alpha i \not{\partial} \nu_L^\alpha - \frac{1}{2} m_\nu^{\alpha\beta} \overline{(\nu_L^c)^\alpha} \nu_L^\beta - \frac{1}{2} y^{\alpha\beta} \phi \overline{(\nu_L^c)^\alpha} \nu_L^\beta.$$

The interaction term can be generated in a gauge invariant way by coupling ϕ to heavy right-handed neutrinos N_R (introduced in seesaw type-I)

- ▶ we assume $y = y_0(m_\nu/0.1\text{eV})$

$$\text{Vector } \mathcal{L}_{\text{vector}} = \bar{\nu}_L^\alpha i \not{\partial} \nu_L^\alpha - \frac{1}{2} m_\nu^{\alpha\beta} \overline{(\nu_L^c)^\alpha} \nu_L^\beta + g Q^{\alpha\beta} \phi^\mu \bar{\nu}_L^\alpha \gamma_\mu \nu_L^\beta.$$

- ▶ ϕ^μ as the $L_\mu - L_\tau$ symmetry gauge boson with couplings $Q^{\alpha\beta} = \text{diag}(0, 1, -1)$
- ▶ if $L_\mu - L_\tau$ breaking occurs at TeV scale, with $m_\phi \sim 10^{-22}$ eV we require coupling $g \sim 10^{-30}$ which can be probed.

Model II

- alternatively, ϕ^μ could couple to the SM via mixing with a much heavier $L_\mu - L_\tau$ gauge boson K^μ (term $\epsilon\phi^{\mu\nu}K_{\mu\nu}$)

$$\mathcal{L}_{\mu-\tau} = -\frac{1}{4}K_{\mu\nu}K^{\mu\nu} + \bar{L}^\alpha (i\not{\partial} + g_{\mu-\tau}Q_{\mu-\tau}^\alpha\gamma_\mu K^\mu)L^\alpha + \bar{e}_R^\alpha (i\not{\partial} + g_{\mu-\tau}Q_{\mu-\tau}^\alpha\gamma_\mu K^\mu)e_R^\alpha$$

$$+ (D^\mu S)^\dagger(D_\mu S) + \mu_S^2 S^\dagger S - \lambda_S(S^\dagger S)^2$$

$$\mathcal{L}_{\text{dark}} = -\frac{1}{4}\phi_{\mu\nu}\phi^{\mu\nu} + \frac{1}{2}\epsilon\phi_{\mu\nu}K^{\mu\nu} + \frac{1}{2}(\partial_\mu\sigma + m_1\phi_\mu + m_2K_\mu)^2$$

kinetic and mass term in matrix form : $V = (\phi, K)^T$

$$\mathcal{L} \supset -\frac{1}{4}V_{\mu\nu}^T \begin{pmatrix} 1 & -\epsilon \\ -\epsilon & 1 \end{pmatrix} V^{\mu\nu} + \frac{1}{2}V_\mu^T \begin{pmatrix} m_1^2 & m_1 m_2 \\ m_1 m_2 & m_2^2 + (g_{\mu-\tau}v_S)^2 \end{pmatrix} V^\mu$$

after two unitary transformations

$$\begin{pmatrix} \phi \\ K \end{pmatrix} = U \begin{pmatrix} \tilde{\phi} \\ \tilde{K} \end{pmatrix} \equiv U_1 U_2 \begin{pmatrix} \tilde{\phi} \\ \tilde{K} \end{pmatrix}$$

we identify gauge boson masses and the effective coupling $y_i = \frac{m_i}{g_{\mu-\tau}v_S}$

$$\mathcal{L}_{\text{int}} = \left(-\frac{y_1^2\epsilon}{1-y_1^2} - \frac{y_1 y_2}{1-y_1^2} \right) g_{\mu-\tau} Q_{\mu-\tau}^\alpha \tilde{\phi}^\mu (\bar{L}^\alpha \gamma_\mu L^\alpha + \bar{e}_R^\alpha \gamma_\mu e_R^\alpha)$$

- ▶ neutrino masses are generated by introducing 3 RH neutrinos with the following charges under $SU(3)_c \times SU(2)_L \times U(1)_Y \times U(1)_{L_\mu-L_\tau}$

$$N_1 \sim (1, 1, 0)(0), \quad N_2 \sim (1, 1, 0)(+1), \quad N_3 \sim (1, 1, 0)(-1)$$

$$\mathcal{L}_{yuk} = \frac{1}{2} a \bar{N}_1^c N_1 + \frac{1}{2} b (\bar{N}_2^c N_3 + \bar{N}_3^c N_2) + \lambda_e \bar{L}_e \tilde{H} N_1 + \lambda_\mu \bar{L}_\mu \tilde{H} N_2 + \lambda_\tau \bar{L}_\tau \tilde{H} N_3 + h.c. + \lambda_S^{12} \bar{N}_1^c N_2 S + \lambda_S^{13} \bar{N}_1^c N_3 S^* + h.c.$$

$$m_D = \begin{pmatrix} m_{\nu_e} & 0 & 0 \\ 0 & m_{\nu_\mu} & 0 \\ 0 & 0 & m_{\nu_\tau} \end{pmatrix}, \quad m_R = \begin{pmatrix} a & s & t \\ s & 0 & b \\ t & b & 0 \end{pmatrix},$$
$$m_{\nu_j} \equiv \lambda_j v / \sqrt{2}, \quad s \equiv \lambda_S^{12} v_X \quad \text{and} \quad t \equiv \lambda_S^{13} v_X.$$

$$m_\nu \simeq -m_D \cdot m_R^{-1} \cdot m_D.$$

MSW Potential

- ▶ Coherent Forward Scattering of Neutrinos on Fuzzy DM
- ▶ scalar DM


$$V_{\text{eff}} = \frac{1}{2E_\nu} \left(\phi (y m_\nu + m_\nu y) + \phi^2 y^2 \right), \quad \phi = \frac{\sqrt{2\rho_\phi}}{m_\phi} \cos(m_\phi t),$$

- ▶ vector DM

$$V_{\text{eff}} = -\frac{1}{2E_\nu} \left(2(p_\nu \cdot \phi)gQ + g^2 Q^2 \phi^2 \right). \quad \phi^\mu = \frac{\sqrt{2\rho_\phi}}{m_\phi} \xi^\mu \cos(m_\phi t).$$


- ▶ $V_{\mu\mu}^{(T,U)} = V_{\tau\tau}^{(T,U)} = \frac{g^2 \rho_\phi}{E_\nu m_\phi^2} \cos^2(m_\phi t)$

- ▶ for polarized DM we evaluate $p_\nu \cdot \phi$ assuming the polarization axis to be parallel to the ecliptic plane



Methods

- ▶ We have implemented the potential in GLoBES [Huber et al. 0701187,0407333](#)
- ▶ the time dependence of matter potential induces time dependent oscillation probabilities
- ▶ we evaluate the oscillation probabilities at several fixed times and interpolate using a second order polynomial in $\cos(m_\phi t)$




$$P(\nu_\alpha^- \rightarrow \nu_\beta^-) = P_0^{\alpha\beta}(E_\nu) + P_1^{\alpha\beta}(E_\nu) \cdot V(t) + P_2^{\alpha\beta}(E_\nu) \cdot V(t)^2 + \dots$$

- ▶ the probability is then averaged in a given time interval T


$$\bar{P}(E) = \frac{1}{T} \int_0^T dt P(E_\nu, t)$$

Constraints



- ▶ for vector DM, the sensitivity is more than ten orders of magnitude better in the polarized case
- ▶ for scalar and polarized vector DM acceleration-based experiments give stronger limits and sensitivities
- ▶ for unpolarized vector DM, experiments at lower energies are better (energy dependence of the potential)

Impact on Solar and Astrophysical neutrinos



Solar


- ▶ adiabatic evolution in the sun Sun
- ▶ survival probability of electron flavor

$$P_{ee}(E_\nu) = \sum_i |U_{ei}^\ominus|^2 |U_{ei}^\oplus|^2$$
- ▶ fitted data from Borexino, Super-K and SNO

Astrophysical

- ▶ obtaining constraints from optical depth $\tau_\nu(E_\nu) = \sigma_{\nu\phi}(E_\nu) X_\phi m_\phi^{-1}$ with

$$X_\phi \equiv \int_{l.o.s} dl \rho_\phi$$
- ▶ much weaker limits in comparison to oscillation exp.




Summary




- ▶ fuzzy DM is an interesting alternative to WIMP
- ▶ fuzzy neutrinophilic DM has recently received attention (Berlin 1608.01307, Krnjaić et al. 1705.06740)
- ▶ we have demonstrated that unique opportunities exist at current and future neutrino oscillation experiments to probe interactions between neutrinos and ultra-light DM particles
- ▶ possible connections with LHCb anomalies

BACKUP



BSM neutrino oscillation physics: eV-scale ν_s



Reactor antineutrino fluxes: status and challenges (Neutrino 2016)
Huber 2011, Mueller et al. 2011



BSM neutrino oscillation physics: eV-scale ν_s



- ▶ tension between appearance signals and disappearance data (see for instance global fits – Kopp et al, Giunti et al., Conrad et al.)
- ▶ experiments such as MicroBooNE and SBL reactor experiments (STEREO...) plan to address the anomaly

BSM neutrino oscillation physics:

“Standard” Non-Standard Interactions

$$i \frac{d}{dt} \begin{pmatrix} \nu_e \\ \nu_\mu \\ \nu_\tau \end{pmatrix} = \frac{1}{2E} \left[U \begin{pmatrix} 0 & 0 & 0 \\ 0 & \Delta m_{21}^2 & 0 \\ 0 & 0 & \Delta m_{31}^2 \end{pmatrix} U^\dagger + 2\sqrt{2}EG_F n_e \begin{pmatrix} 1 + \epsilon_{ee} & \epsilon_{e\mu} & \epsilon_{e\tau} \\ \epsilon_{e\mu}^* & \epsilon_{\mu\mu} & \epsilon_{\mu\tau} \\ \epsilon_{e\tau}^* & \epsilon_{\mu\tau}^* & \epsilon_{\tau\tau} \end{pmatrix} \right] \begin{pmatrix} \nu_e \\ \nu_\mu \\ \nu_\tau \end{pmatrix}$$

