

Going global

Combining electroweak precision, diboson,
Higgs, and top data to search for new physics

Ken Mimasu

King's College London

High energy and nuclear physics seminar, Uppsala University

22nd April 2021

[J. Ellis, M. Madigan, KM, V. Sanz & T. You; arXiv:2012.02779]

fitmaker <https://gitlab.com/kenmimasu/fitrepo>


[G. Durieux, C. Degrande, F. Maltoni, KM, C. Zhang, E. Vryonidou; arXiv:2008.11743]

SMEFTatNLO <http://feynrules.irmp.ucl.ac.be/wiki/SMEFTatNLO>

Where are we?

10 years since the start of LHC Run 1

- No clear sign of new physics at the TeV scale
- Direct searches are saturating the energy frontier



[CERN-EP-2020-049]

ATLAS Exotics Searches* - 95% CL Upper Exclusion Limits
Status: May 2020

Model	ℓ, γ	Jets [†]	E_T^{miss}	$\int \mathcal{L} dt [\text{fb}^{-1}]$	Limit	Reference
Extra dimensions						
ADD $G_{KK} + g/q$	0 e, μ	1 – 4 j	Yes	36.1	M_D	1711.03301
ADD non-resonant $\gamma\gamma$	2 γ	–	–	36.7	M_S	1707.04147
ADD QBH	–	2 j	–	37.0	M_{bh}	1703.09127
ADD BH high Σp_T	$\geq 1 e, \mu$	$\geq 2 j$	–	3.2	M_{bh}	1606.02265
ADD BH multijet	–	$\geq 3 j$	–	3.6	M_{bh}	1512.02586
RS1 $G_{KK} \rightarrow \gamma\gamma$	2 γ	–	–	36.7	G_{KK} mass	1707.04147
Bulk RS $G_{KK} \rightarrow WW/ZZ$	multi-channel	–	–	36.1	G_{KK} mass	1808.02380
Bulk RS $G_{KK} \rightarrow WV \rightarrow \ell\nu qq$	1 e, μ	2 j / 1 J	Yes	139	G_{KK} mass	2004.14636
Bulk RS $G_{KK} \rightarrow tt$	1 e, μ	$\geq 1 b, \geq 1 J/2j$	Yes	36.1	g_{KK} mass	1804.10823
2UED / RPP	1 e, μ	$\geq 2 b, \geq 3 j$	Yes	36.1	KK mass	1803.09678
Gauge bosons						
SSM $Z' \rightarrow \ell\ell$	2 e, μ	–	–	139	Z' mass	1903.06248
SSM $Z' \rightarrow \tau\tau$	2 τ	–	–	36.1	Z' mass	1709.07242
Leptophobic $Z' \rightarrow bb$	–	2 b	–	36.1	Z' mass	1805.09299
Leptophobic $Z' \rightarrow tt$	0 e, μ	$\geq 1 b, \geq 2 J$	Yes	139	Z' mass	2005.05138
SSM $W' \rightarrow \ell\nu$	1 e, μ	–	–	139	W' mass	1906.05609
SSM $W' \rightarrow \tau\nu$	1 τ	–	–	36.1	W' mass	1801.06992
HVT $V' \rightarrow WV \rightarrow \ell\nu qq$ model B	1 e, μ	2 j / 1 J	Yes	139	V' mass	2004.14636
HVT $V' \rightarrow WV \rightarrow qqqq$ model B	0 e, μ	2 J	–	36.1	V' mass	1906.08589
HVT $V' \rightarrow WH/ZH$ model B	multi-channel	–	–	36.1	V' mass	1712.06518
LRSM $W_R \rightarrow tb$	0 e, μ	$\geq 1 b, \geq 2 J$	–	139	W_R mass	CERN-EP-2020-073
LRSM $W_R \rightarrow \mu N_R$	2 μ	1 J	–	80	W_R mass	1807.10473
Cl						
Cl $qqqq$	–	2 j	–	37.0	Λ	1703.09127
Cl $\ell\ell qq$	2 e, μ	–	–	139	Λ	1811.02305
Cl $t\bar{t}tt$	$\geq 1 e, \mu$	$\geq 1 b, \geq 1 j$	Yes	36.1	Λ	
DM						
Axial-vector mediator (Dirac DM)	0 e, μ	1 – 4 j	Yes	36.1	m_{min}	1711.03301
Colored scalar mediator (Dirac DM)	0 e, μ	1 – 4 j	Yes	36.1	m_{min}	1711.03301
VV_{XX} EFT (Dirac DM)	0 e, μ	1 J, $\leq 1 j$	Yes	3.2	M_s	1608.02372
Scalar reson. $\phi \rightarrow t\chi$ (Dirac DM)	0-1 e, μ	1 b, 0-1 J	Yes	36.1	m_p	1812.09743
LQ						
Scalar LQ 1 st gen	1.2 e	$\geq 2 j$	Yes	36.1	LQ mass	1902.00377
Scalar LQ 2 nd gen	1.2 μ	$\geq 2 j$	Yes	36.1	LQ mass	1902.00377
Scalar LQ 3 rd gen	2 τ	2 b	–	36.1	LQ ₃ mass	1902.08103
Scalar LQ 3 rd gen	0-1 e, μ	2 b	Yes	36.1	LQ ₃ mass	1902.08103
Heavy quarks						
VLQ $T T \rightarrow Ht/Zt/Wb + X$	multi-channel	–	–	36.1	T mass	1808.02343
VLQ $BB \rightarrow Wt/Zb + X$	–	–	–	36.1	B mass	1808.02343
VLQ $T_{5/3} T_{5/3} \rightarrow Wt + X$	$2(S) \geq 3$ e, μ	$\geq 1 b, \geq 1 j$	Yes	36.1	$T_{5/3}$ mass	1807.11883
VLQ $Y \rightarrow Wb + X$	1 e, μ	$\geq 1 b, \geq 1 j$	Yes	36.1	Y mass	1812.07343
VLQ $B \rightarrow Hb + X$	0 e, $\mu, 2 \gamma$	$\geq 1 b, \geq 1 j$	Yes	79.8	B mass	ATLAS-CONF-2018-024
VLQ $QQ \rightarrow WqWq$	1 e, μ	$\geq 4 j$	Yes	20.3	Q mass	1509.04261
Excited fermions						
Excited quark $q^* \rightarrow qg$	–	2 j	–	139	q^* mass	1910.08447
Excited quark $q^* \rightarrow q\gamma$	1 γ	1 j	–	36.7	q^* mass	1709.10440
Excited quark $b^* \rightarrow bg$	–	1 b, 1 j	–	36.1	b^* mass	1805.09299
Excited lepton ℓ^*	3 e, μ	–	–	20.3	ℓ^* mass	1411.2921
Excited lepton ν^*	3 e, μ, τ	–	–	20.3	ν^* mass	1411.2921
Other						
Type III Seesaw	1 e, μ	$\geq 2 j$	Yes	79.8	N^0 mass	ATLAS-CONF-2018-020
LRSM Majorana ν	2 μ	2 j	–	36.1	N_R mass	1809.11105
Higgs triplet $H^{\pm} \rightarrow \ell\ell$	2,3,4 e, μ (SS)	–	–	36.1	H^{\pm} mass	1710.09748
Higgs triplet $H^{\pm} \rightarrow \ell\tau$	3 e, μ, τ	–	–	20.3	400 GeV	1411.2921
Multi-charged particles	–	–	–	36.1	multi-charged particle mass	1812.03673
Magnetic monopoles	–	–	–	34.4	monopole mass	1905.10130

[TeV] : 1 3 5 10

Top, Higgs, Diboson & EW fit to the SMEFT

What have we learnt?

BSM states are too...

Weakly coupled

rate limited

Room for improvement with increasing luminosity

Still 20x more data to come

Exotic

*we aren't looking in
the right place*

Limited by our creativity

Work for theorists & experimentalists to motivate & enable searches for new signatures

Heavy

*kinematically
out of reach*

Worst-case scenario from direct search point of view

Complemented by indirect searches


A tremendous amount about the SM!

- Higgs discovery & properties \Rightarrow precision LHC programme

The LHC explorer


Many new processes observed at the LHC for the first time

Main Higgs production modes




ggF, VH, VBF, ttH

Rare top production



$tttt, ttbb$

Weak boson scattering



VBS, VVV

Each opens a new window, through which we can

- Improve our understanding of the SM
- Search for new physics via new interactions

The SM is broken

Theory & matter content rich with symmetry & structure

$$SU(3)_C \times SU(2)_L \times U(1)_Y$$

$$\varphi = \begin{pmatrix} G^+ \\ v + h + iG^0 \end{pmatrix} : \mathbf{2}_{\frac{1}{2}}$$

Electroweak symmetry breaking

- Offers a **parametrisation**: lacks dynamical origin for the weak scale

Symmetry \leftrightarrow Constraints/Relations

$$y_f \bar{F}_L f_R \varphi \quad (D^\mu \varphi)^\dagger (D_\mu \varphi)$$

Mass \leftrightarrow Higgs coupling

$$\frac{1}{4} W_{\mu\nu}^a W_a^{\mu\nu} \quad i \bar{F} \not{D} F$$

Self-interactions \leftrightarrow Gauge currents

New physics can indirectly perturb this delicate balance

The indirect way



“...the direct method may be used...but indirect methods will be needed in order to secure victory.”

“...there are not more than two methods of attack – the direct and the indirect;... Who can exhaust the possibilities of their combination?”

— Sun Tzu, *The Art of War*


Energy & precision

Paradigm shift at the energy frontier for BSM searches


Direct (bumps)

Indirect (tails)

⇒ New physics is heavy



Heavy new physics
Precision measurements
High energy



**Standard Model
Effective Field Theory
(SMEFT)**

A QFT parameter space for BSM interactions between SM particles

SMEFT: SM v2.0

$$\mathcal{L}_{\text{eff}} = \sum_i \frac{c_i \mathcal{O}_i^D}{\Lambda^{D-4}}$$

SM is low energy effective description

- Supplemented by a tower of irrelevant operators
- Respecting low energy field content & symmetries

$$SU(3)_c \times SU(2)_L \times U(1)_Y$$

$$\varphi = \begin{pmatrix} G^+ \\ v + h + iG^0 \end{pmatrix} : \mathbf{2}_{\frac{1}{2}}$$


aTGC	$X^3 : \epsilon_{IJK} W_{\mu\nu}^I W^{J,\nu\rho} W_{\rho}^{K,\mu}$	$X^2 H^2 : (\varphi^\dagger \varphi)^2 G_{\mu\nu}^a G_a^{\mu\nu}$	ggh(h)
λ_h	$H^6 : (\varphi^\dagger \varphi)^3$	$H^4 D^2 : (\varphi^\dagger D^\mu \varphi)^* (\varphi^\dagger D^\mu \varphi)$	δM_Z
y_f	$\psi^2 H^3 : (\varphi^\dagger \varphi)^2 (\bar{q}_i u_j \tilde{\varphi})$	$\psi^2 X H : (\bar{q}_i \sigma^{\mu\nu} u_j \tilde{\varphi}) B_{\mu\nu}$	'dipole'
ffV	$\psi^2 H^2 D : (\varphi^\dagger \overleftrightarrow{D}_\mu \varphi) (\bar{q}_i \gamma^\mu q_j)$	$\psi^4 : (\bar{q}_i \gamma^\mu q_j) (\bar{q}_k \gamma_\mu q_l)$	4F

More than 'just' a parametrisation of ignorance

- Unlike anomalous couplings
- Finite energy range ($\sim \Lambda$)
- Renormalisable QFT (order-by-order)
- Well defined matching procedure

SMEFT strategy

$$\mathcal{L}_{\text{EFT}} = \mathcal{L}_{\text{SM}} + \sum_i \frac{c_i^{(6)}}{\Lambda^2} O_i^{(6)} + \mathcal{O}(\Lambda^{-4})$$



Map coefficients to the data once and for all

SMEFT is a way to test many BSM scenarios

- Economical
- Well developed

UV matching quasi-automated

- Tree-level dictionary
[de Blas et al.; JHEP 03 (2018) 109]
- Universal one loop effective action
[Henning, Lu & Murayama; JHEP 01 (2016) 023]
[Drozd et al.; JHEP 03 (2016) 180]

RGE are known

- [Alonso*, Jenkins, Manohar & Trott;
JHEP 1310 (2013) 087,
JHEP 1401 (2014) 035
JHEP 1404 (2014) 159*]

Mature MC tools

SMEFTsim, SMEFTatNLO, dim6top,...

SMEFT is...

$$\mathcal{L}_{\text{eff}} = \sum_i \frac{c_i \mathcal{O}_i^D}{\Lambda^{D-4}}$$

Model independent

- Underlying assumptions

Heavy new physics: $M > E_{\text{exp}}$

SM field content & gauge symmetries

Linear EWSB: Higgs = doublet

Systematically improvable

- Double expansion

higher dim. $\frac{E^2}{\Lambda^2}$ & $\{g_s, g, g'\}$ more loops

Global

- Model independence: we don't know what operators NP will generate
- Patterns & correlations among observables are key
- Ultimate goal: complete SMEFT likelihood confronted with HEP data

EWPO, Higgs, multiboson, top, DY, flavor, ...

Established part of LHC programme

SMEFT interpretation

Ingredients:

$$\Delta o_n = o_n^{\text{EXP}} - o_n^{\text{SM}} = \sum_i \frac{a_{n,i}^{(6)}(\mu) c_i^{(6)}(\mu)}{\Lambda^2} + \mathcal{O}\left(\frac{1}{\Lambda^3}\right)$$

Global nature

As many observables as possible

Identify patterns & correlations in fits

Exploit energy-growth

Sensitivity

Experiment:
Best measurements & understanding of uncertainties and correlations

Theory:
Best available predictions for observables (NLO, NNLO, N3LO, ...)

Interpretation

Relies on accurate knowledge of the size & correlation among a_i

Determining $c_i^{(6)}$ requires most precise available SMEFT predictions

Status in a nutshell

Global new physics searches via high precision/energy

- Z & W-pole data: handle on the EW gauge sector [Han & Skiba; PRD 71 (2005) 075009]
[Falkowski & Riva; JHEP 02 (2015) 039]
- LHC: thriving Higgs & top programmes
- Probing gauge interactions at high energy (VV, VBS, VVV, ...)

How much cross-talk? Where does being global matter?

We know that Higgs data greatly complements LEP

- Access unconstrained directions in parameter space
- Allows for a closed fit to flavor-universal SMEFT
- Crucial to combine EWPO, Diboson & Higgs data

[Corbett et al.; PRD 87 (2013) 015022]


[Pomarol & Riva; JHEP 01 (2014) 151]

[Ellis, Sanz & You; JHEP 03 (2015) 157]

[Biekötter Corbett & Plehn; SciPost Phys 6 (2019) 6, 064]...

[Ellis et al.; JHEP 06

(2018) 146]



Top & Higgs


Inextricably linked in the SM

- Yukawa interaction controls ggF
- Strong BSM motivation to study tops

ggF is well measured now


- Does not exclude top partners, anomalous Yukawa!

C_{HG} Point-like
 C_{tH} Yukawa
 C_{tG} Dipole




Need more data to break degeneracy

- $t\bar{t}H$ production for direct Yukawa measurement
- $t\bar{t}$ data to constrain dipole




Blind direction in BSM scenarios
Effective coupling \equiv degeneracy



The role of top data


$t\bar{t}$ cross section measurements constrain C_{tG}

- Indirectly improve bounds on C_{HG} and C_{tH}



Several other new interactions can affect $t\bar{t}$

- Notably $q\bar{q}t\bar{t}$ operators, of which there are many (14)
- To what extent do these limit ultimate NP sensitivity in top/Higgs sector?



Can only be addressed in combined fit

- Beyond tree-level (at least for ggF) *[Degrade et al.; arXiv:2008.11743]*
<http://feynrules.irmp.ucl.ac.be/wiki/SMEFTatNLO>
- Identify other cross-talk (non-trivial correlations)
- Broaden range of applicability to UV models

The fit

arXiv:2012.02779 **Top, Higgs, Diboson and Electroweak Fit to the Standard Model Effective Field Theory**

John Ellis,^{a,b,c} Maeve Madigan,^d Ken Mimasu,^a Veronica Sanz^{e,f} and Tevong You^{b,d,g}

Global SMEFT interpretation of 4 categories of data

- 14 • Electroweak Precision Observables (EWPO): Z-pole & W-mass [Ellis et al.; JHEP 06 (2018) 146]
- 118 • LEP2 & LHC diboson production: differential WW, WZ, Zjj
- 72 • Higgs measurements: signal strengths & STXS
- 137 • Top data: single-top, ttbar & asymmetries, ttV, tZ, tW

Based on

Big thanks to authors of
SMEFiT analysis
[JHEP 04 (2019) 100]
for sharing some of their
top predictions

341 measurements across categories

- Chosen to be statistically independent & maximise reach
- Correlations included when publicly available (mostly are)

Linear EFT approximation: $\mu_X \equiv \frac{X}{X_{SM}} = 1 + \sum_i a_i^X \frac{C_i}{\Lambda^2} + \mathcal{O}\left(\frac{1}{\Lambda^4}\right)$

Theory

[Grzadkowski et al.; JHEP 10 (2010) 085]

X^3		H^6 and H^4D^2		$\psi^2 H^3$	
\mathcal{O}_G	$f^{ABC} G_\mu^{A\nu} G_\nu^{B\rho} G_\rho^{C\mu}$	\mathcal{O}_H	$(H^\dagger H)^3$	\mathcal{O}_{eH}	$(H^\dagger H)(\bar{l}_p e_r H)$
$\mathcal{O}_{\tilde{G}}$	$f^{ABC} \tilde{G}_\mu^{A\nu} G_\nu^{B\rho} G_\rho^{C\mu}$	$\mathcal{O}_{H\square}$	$(H^\dagger H)\square(H^\dagger H)$	\mathcal{O}_{uH}	$(H^\dagger H)(\bar{q}_p u_r \tilde{H})$
\mathcal{O}_W	$\varepsilon^{IJK} W_\mu^{I\nu} W_\nu^{J\rho} W_\rho^{K\mu}$	\mathcal{O}_{HD}	$(H^\dagger D^\mu H)^*(H^\dagger D_\mu H)$	\mathcal{O}_{dH}	$(H^\dagger H)(\bar{q}_p d_r H)$
$\mathcal{O}_{\tilde{W}}$	$\varepsilon^{IJK} \tilde{W}_\mu^{I\nu} W_\nu^{J\rho} W_\rho^{K\mu}$				
$X^2 H^2$		$\psi^2 XH$		$\psi^2 H^2 D$	
\mathcal{O}_{HG}	$H^\dagger H G_{\mu\nu}^A G^{A\mu\nu}$	\mathcal{O}_{eW}	$(\bar{l}_p \sigma^{\mu\nu} e_r) \tau^I H W_{\mu\nu}^I$	$\mathcal{O}_{Hl}^{(1)}$	$(H^\dagger i \overset{\leftrightarrow}{D}_\mu H)(\bar{l}_p \gamma^\mu l_r)$
$\mathcal{O}_{H\tilde{G}}$	$H^\dagger H \tilde{G}_{\mu\nu}^A G^{A\mu\nu}$	\mathcal{O}_{eB}	$(\bar{l}_p \sigma^{\mu\nu} e_r) H B_{\mu\nu}$	$\mathcal{O}_{Hl}^{(3)}$	$(H^\dagger i \overset{\leftrightarrow}{D}_\mu^I H)(\bar{l}_p \tau^I \gamma^\mu l_r)$
\mathcal{O}_{HW}	$H^\dagger H W_{\mu\nu}^I W^{I\mu\nu}$	\mathcal{O}_{uG}	$(\bar{q}_p \sigma^{\mu\nu} T^A u_r) \tilde{H} G_{\mu\nu}^A$	\mathcal{O}_{He}	$(H^\dagger i \overset{\leftrightarrow}{D}_\mu H)(\bar{e}_p \gamma^\mu e_r)$
$\mathcal{O}_{H\tilde{W}}$	$H^\dagger H \tilde{W}_{\mu\nu}^I W^{I\mu\nu}$	\mathcal{O}_{uW}	$(\bar{q}_p \sigma^{\mu\nu} u_r) \tau^I \tilde{H} W_{\mu\nu}^I$	$\mathcal{O}_{Hq}^{(1)}$	$(H^\dagger i \overset{\leftrightarrow}{D}_\mu H)(\bar{q}_p \gamma^\mu q_r)$
\mathcal{O}_{HB}	$H^\dagger H B_{\mu\nu} B^{\mu\nu}$	\mathcal{O}_{uB}	$(\bar{q}_p \sigma^{\mu\nu} u_r) \tilde{H} B_{\mu\nu}$	$\mathcal{O}_{Hq}^{(3)}$	$(H^\dagger i \overset{\leftrightarrow}{D}_\mu^I H)(\bar{q}_p \tau^I \gamma^\mu q_r)$
$\mathcal{O}_{H\tilde{B}}$	$H^\dagger H \tilde{B}_{\mu\nu} B^{\mu\nu}$	\mathcal{O}_{dG}	$(\bar{q}_p \sigma^{\mu\nu} T^A d_r) H G_{\mu\nu}^A$	\mathcal{O}_{Hu}	$(H^\dagger i \overset{\leftrightarrow}{D}_\mu H)(\bar{u}_p \gamma^\mu u_r)$
\mathcal{O}_{HWB}	$H^\dagger \tau^I H W_{\mu\nu}^I B^{\mu\nu}$	\mathcal{O}_{dW}	$(\bar{q}_p \sigma^{\mu\nu} d_r) \tau^I H W_{\mu\nu}^I$	\mathcal{O}_{Hd}	$(H^\dagger i \overset{\leftrightarrow}{D}_\mu H)(\bar{d}_p \gamma^\mu d_r)$
$\mathcal{O}_{H\tilde{W}B}$	$H^\dagger \tau^I H \tilde{W}_{\mu\nu}^I B^{\mu\nu}$	\mathcal{O}_{dB}	$(\bar{q}_p \sigma^{\mu\nu} d_r) H B_{\mu\nu}$	\mathcal{O}_{Hud}	$i(\tilde{H}^\dagger D_\mu H)(\bar{u}_p \gamma^\mu d_r)$

$(\bar{L}L)(\bar{L}L)$		$(\bar{R}R)(\bar{R}R)$		$(\bar{L}L)(\bar{R}R)$	
\mathcal{O}_{ll}	$(\bar{l}_p \gamma_\mu l_r)(\bar{l}_s \gamma^\mu l_t)$	\mathcal{O}_{ee}	$(\bar{e}_p \gamma_\mu e_r)(\bar{e}_s \gamma^\mu e_t)$	\mathcal{O}_{le}	$(\bar{l}_p \gamma_\mu l_r)(\bar{e}_s \gamma^\mu e_t)$
$\mathcal{O}_{qq}^{(1)}$	$(\bar{q}_p \gamma_\mu q_r)(\bar{q}_s \gamma^\mu q_t)$	\mathcal{O}_{uu}	$(\bar{u}_p \gamma_\mu u_r)(\bar{u}_s \gamma^\mu u_t)$	\mathcal{O}_{lu}	$(\bar{l}_p \gamma_\mu l_r)(\bar{u}_s \gamma^\mu u_t)$
$\mathcal{O}_{qq}^{(3)}$	$(\bar{q}_p \gamma_\mu \tau^I q_r)(\bar{q}_s \gamma^\mu \tau^I q_t)$	\mathcal{O}_{dd}	$(\bar{d}_p \gamma_\mu d_r)(\bar{d}_s \gamma^\mu d_t)$	\mathcal{O}_{ld}	$(\bar{l}_p \gamma_\mu l_r)(\bar{d}_s \gamma^\mu d_t)$
$\mathcal{O}_{lq}^{(1)}$	$(\bar{l}_p \gamma_\mu l_r)(\bar{q}_s \gamma^\mu q_t)$	\mathcal{O}_{eu}	$(\bar{e}_p \gamma_\mu e_r)(\bar{u}_s \gamma^\mu u_t)$	\mathcal{O}_{qe}	$(\bar{q}_p \gamma_\mu q_r)(\bar{e}_s \gamma^\mu e_t)$
$\mathcal{O}_{lq}^{(3)}$	$(\bar{l}_p \gamma_\mu \tau^I l_r)(\bar{q}_s \gamma^\mu \tau^I q_t)$				$(\bar{q}_p \gamma_\mu q_r)(\bar{u}_s \gamma^\mu u_t)$
$(\bar{L}R)(\bar{R}L)$ and $(\bar{L}R)(\bar{L}R)$		B-violating			
\mathcal{O}_{ledq}	$(\bar{l}_p^j e_r)(\bar{d}_s q_t^j)$	\mathcal{O}_{duq}	$\varepsilon^{\alpha\beta\gamma} \varepsilon_{jk} [(d_p^\alpha)^T C u_r^\beta] [(q_s^\gamma)^T C l_t^k]$		
\mathcal{O}_{quqd}	$(\bar{q}_p^j u_r) \varepsilon_{jk} (\bar{q}_s^k d_t)$	\mathcal{O}_{qqu}	$\varepsilon^{\alpha\beta\gamma} \varepsilon_{jk} [(q_p^\alpha)^T C q_r^\beta k] [(u_s^\gamma)^T C e_t]$		
$\mathcal{O}_{quqd}^{(8)}$	$(\bar{q}_p^j T^A u_r) \varepsilon_{jk} (\bar{q}_s^k T^A d_t)$	\mathcal{O}_{qqq}	$\varepsilon^{\alpha\beta\gamma} \varepsilon_{jn} \varepsilon_{km} [(q_p^\alpha)^T C q_r^\beta k] [(q_s^\gamma)^T C l_t^n]$		
$\mathcal{O}_{lequ}^{(1)}$	$(\bar{l}_p^j e_r) \varepsilon_{jk} (\bar{q}_s^k u_t)$	\mathcal{O}_{duu}	$\varepsilon^{\alpha\beta\gamma} [(d_p^\alpha)^T C u_r^\beta] [(u_s^\gamma)^T C e_t]$		
$\mathcal{O}_{lequ}^{(3)}$	$(\bar{l}_p^j \sigma_{\mu\nu} e_r) \varepsilon_{jk} (\bar{q}_s^k \sigma^{\mu\nu} u_t)$				

Warsaw basis with CP & B conservation

- Full ‘bosonic’ sector: Higgs, triple-gauge & gauge-Higgs
- Scenario 1: Flavor-**universal** degrees of freedom

$$U(3)_L \times U(3)_e \times U(3)_Q \times U(3)_u \times U(3)_d + \text{Yukawas: } \mathcal{O}_{tH}, \mathcal{O}_{bH}, \mathcal{O}_{\tau H}, \mathcal{O}_{\mu H}$$

- Scenario 2: **top**-centric flavor symmetry

$$U(3)_L \times U(3)_e \times U(2)_Q \times U(2)_u \times U(3)_d$$



cf. Minimal flavor violation

[Buras et al.; PLB 500 (2001) 161]

[D'Ambrosio et al.; NPB 645 (2002) 155]

[Aguilar-Saavedra et al.; arXiv:1802.07237]

Degrees of freedom

EWPO:	$\mathcal{O}_{HWB}, \mathcal{O}_{HD}, \mathcal{O}_{ll}, \mathcal{O}_{Hl}^{(3)}, \mathcal{O}_{Hl}^{(1)}, \mathcal{O}_{He}, \mathcal{O}_{Hq}^{(3)}, \mathcal{O}_{Hq}^{(1)}, \mathcal{O}_{Hd}, \mathcal{O}_{Hu},$	
Bosonic:	$\mathcal{O}_{H\square}, \mathcal{O}_{HG}, \mathcal{O}_{HW}, \mathcal{O}_{HB}, \mathcal{O}_W, \mathcal{O}_G,$	
Yukawa:	$\mathcal{O}_{\tau H}, \mathcal{O}_{\mu H}, \mathcal{O}_{b H}, \mathcal{O}_{t H},$	20
Top 2F:	$\mathcal{O}_{HQ}^{(3)}, \mathcal{O}_{HQ}^{(1)}, \mathcal{O}_{Ht}, \mathcal{O}_{tG}, \mathcal{O}_{tW}, \mathcal{O}_{tB},$	
Top 4F:	$\mathcal{O}_{Qq}^{3,1}, \mathcal{O}_{Qq}^{3,8}, \mathcal{O}_{Qq}^{1,8}, \mathcal{O}_{Qu}^8, \mathcal{O}_{Qd}^8, \mathcal{O}_{tQ}^8, \mathcal{O}_{tu}^8, \mathcal{O}_{td}^8.$	+ 14

In total: 20(34) d.o.f. for the two flavor scenarios


Dim6top conventions: [Aguilar-Saavedra et al.; arXiv:1802.07237]

Dictated by flavor symmetry & sensitivity of dataset

Linear EFT fit: precludes sensitivity to some ops

- Those that cannot interfere due to helicity/symmetries
- e.g. neutral colour-singlet top 4F operators: $(\bar{q}\gamma^\mu q)(\bar{t}\gamma^\mu t)$ (x 6)
- Four-heavy quark operators in 4top & ttbb (quadratic dominated)

Interplay



Technical details

$$\mu_X \equiv \frac{X}{X_{SM}} = 1 + \sum_i a_i^X \frac{C_i}{\Lambda^2} + \mathcal{O}\left(\frac{1}{\Lambda^4}\right)$$

Exp. data: [HEPdata](#), [WebPlotDigitizer](#), ...

- Construct ‘signal strength’, w.r.t. SM prediction from exp. paper
- Otherwise computed with **MG5**, **fastnlo**, directly from theory papers
- Combine all sources of uncertainty in quadrature (stat., syst., th.)

Theory predictions: **MG5** (**SMEFTsim** & **SMEFTatNLO**)

- LO, parton-level, linear dependence in (α, G_F, M_Z) scheme
- Tree-level + 1-loop gluon fusion Higgs production
- a_i : Effects from production, decays, total width
- No theory error from EFT, assume SM error dominant

The code

fitmaker <https://gitlab.com/kenmimasu/fitrepo>
public-friendly version w/ example notebooks in progress

Main analysis: linearised least-squares fit

$$\chi^2(C_i) = (\vec{y} - \vec{\mu}(C_i))^T \mathbf{V}^{-1} (\vec{y} - \vec{\mu}(C_i)) \quad \mu_\alpha(C_i) = \mu_\alpha^{\text{SM}} + \mathbf{H}_{\alpha i} C_i$$

Best fit $\hat{\vec{C}} = (\mathbf{H}^T \mathbf{V}^{-1} \mathbf{H})^{-1} \mathbf{H}^T \mathbf{V}^{-1} (\vec{y} - \vec{\mu}^{\text{SM}}) \equiv \mathbf{F}^{-1} \vec{\omega}$

$$\mathbf{F} \equiv \mathbf{H}^T \mathbf{V}^{-1} \mathbf{H} \quad , \quad \vec{\omega} \equiv \mathbf{H}^T \mathbf{V}^{-1} (\vec{y} - \vec{\mu}^{\text{SM}}) ,$$

Fisher information

$\mathbf{F}^{-1} \equiv \mathbf{U}$ Covariance matrix of least-squares estimator

$(\chi_{\text{SM}}^2, \hat{\vec{C}}, \mathbf{U})$ fully characterise likelihood

- Individual, profiled/marginalised bounds & correlations
- Principal component analysis (eigensystem of F)

Implemented as part
of the `fitmaker`
framework

Also nested sampling routine for general likelihoods

The code

fitmaker <https://gitlab.com/kenmimasu/fitrepo>
public-friendly version w/ example notebooks in progress

Database of input measurements encoded in `.json` format

- Values, errors, metadata,...

Python-class based definition of theoretical models

- Predictions for observables can be hard-coded
- ...or read-in from a `.json` file

$$\mu_{H \rightarrow 4\ell}^{ggF} = 0.98^{+0.12}_{-0.11}$$

```
{-  
  "observable_name": "mu_ggF_H_ZZ_13",  
  "measurement_name": "mu_ggF_H_ZZ_CMS_Run2",  
  "CDS": "http://cds.cern.ch/record/2706103",  
  "reportnumber": "CMS-PAS-HIG-19-005",  
  "DOI": "",  
  "date": "2020/01/10",  
  "experiment": "CERN-LHC-Run-2, CMS",  
  "description": "Higgs boson signal strength for  
  "value": 0.98,  
  "uncertainty": {  
    "tot": [0.12, 0.11]  
  },  
  "uncertainty_sigma": 1,  
  "th_flat": true  
}-
```

$$\begin{aligned}\mu^{ggF} = & 1 + 35.8C_{HG} - 0.122C_{tH} \\ & - 0.959C_{tG} - 0.121C_{H\square} + \dots\end{aligned}$$

```
{-  
  "observable": "ggF0j",  
  "params": ["CHG", "CuH", "CuG", "CHbox"],  
  "constant": 1.0,  
  "linear": [35.8, -0.122, 0.959, -0.121],  
  "quadratic": [  
    [321.0, -1.095, 8.45, -1.085],  
    [-1.095, 0.00371, -0.02925, 0.003695],  
    [8.45, -0.02925, 0.23, -0.0291],  
    [-1.085, 0.003695, -0.0291, 0.00367]  
  ],  
  "lambda_gen": 1000.0  
}-
```

SMEFT@NLO

Loops & SMEFT: active field in recent years

- Non-universal K-factors in EFT space \Leftrightarrow new information at NLO
- Loop-induced sensitivity (e.g. $gg \rightarrow H$)
- Control theoretical uncertainties
- Experimental interest in higher precision for SMEFT analyses/interpretations

Challenge: many processes \times many operators

- LO \Rightarrow NLO = more cross-talk/operators/complexity
- Automated tools for fixed-order/NLO+PS are essential to the LHC programme

Solution: SMEFT@NLO

- UFO model for `MadGraph5_aMC@NLO`
- Process-independent implementation: SMEFT in top-specific flavor limit

Standard Model Effective Theory at One-Loop in QCD

Céline Degrande, Gauthier Durieux, Fabio Maltoni, Ken Mimasu, Eleni Vryonidou & Cen Zhang, [arXiv:2008.11743](#)

The implementation is based on the Warsaw basis of dimension-six SMEFT operators, after canonical normalization. Electroweak input parameters are taken to be G_F , M_Z , M_W . The CKM matrix is approximated as a unit matrix, and a $U(2)_q \times U(2)_u \times U(3)_d \times (U(1)_l \times U(1)_e)^3$ flavor symmetry is enforced. It forbids all fermion masses and Yukawa couplings except that of the top quark. The model therefore implements the five-flavor scheme for PDFs.

A new coupling order, **NP=2**, is assigned to SMEFT interactions. The cutoff scale **Lambda** takes a default value of 1 TeV^{-2} and can be modified along with the Wilson coefficients in the **param_card**. Operators definitions, normalisations and coefficient names in the UFO model are specified in [definitions.pdf](#). The notations and normalizations of top-quark operator coefficients comply with the LHC TOP WG standards of [1802.07237](#). Note however that the flavor symmetry enforced here is slightly more restrictive than the baseline assumption there (see the [dim6top page](#) for more information). This model has been validated at tree level against the **dim6top** implementation (see [1906.12310](#) and the [comparison details](#)).

Current implementation

UFO model: [SMEFTatNLO_v1.0.tar.gz](#)

The current implementation imposes CP conservation. In the quark sector, it focuses primarily on top-quark interactions. The light-quark current operator, $qq\bar{H}D\bar{H}$, $uu\bar{H}D\bar{H}$, $dd\bar{H}D\bar{H}$, with coefficients **cpq3i**, **cpqMi**, **cpu**, **cpd** are however included. The triple-gluon operator, with coefficient **cG**, is currently not available (see the loop-capable **GGG** implementation). Vertices including more than four scalars or four leptons are not included. Scalar and tensor **QQ11** operators, with coefficients **ct1S3**, **ct1T3**, and **cb1S3**, break our flavor symmetry assumption and are not available for one-loop computations. Top-quark flavor-changing interactions, not compatible with the imposed flavor symmetry, are not included (see the loop-capable [TopFCNC](#) implementation).

Unlike prescribed by the LHC TOP WG, the top quark chromomagnetic-dipole operator coefficient **ctG** is normalized with a factor of the strong coupling, g_S . This normalization factor temporarily ensures compatibility with the 2.X.X series of MadGraph5_aMC@NLO but may be dropped in the future. As with every other appearance of this coupling in MadGraph5_aMC@NLO, its value is renormalisation-group evolved to the QCD renormalization scale (set in the **run_card**).

MG5_aMC>import model SMEFTatNLO

MG5_aMC>generate p p > t t~ NP=2 [QCD]


MG5_aMC>output

MG5_aMC>launch

SMEFT@NLO in STXS

Gluon fusion Simplified Template Cross Sections bins

- LO in the SM is one-loop
- Tree-EFT \times loop-SM + loop-EFT \times loop-SM interference terms
- Heavy top limit is OK for 0-jet, breaks down at high- p_T



Results roadmap

1. Flavor universal: EWPO + diboson + Higgs

2. Top only: EWPO + top

Interlude: Top-Higgs interplay


3. Top-specific : EWPO + diboson + Higgs + top

$U(3)^5$
↓
 $U(2)^2 \times U(3)^3$

EWPO:	$\mathcal{O}_{HWB}, \mathcal{O}_{HD}, \mathcal{O}_{ll}, \mathcal{O}_{Hl}^{(3)}, \mathcal{O}_{Hl}^{(1)}, \mathcal{O}_{He}, \mathcal{O}_{Hq}^{(3)}, \mathcal{O}_{Hq}^{(1)}, \mathcal{O}_{Hd}, \mathcal{O}_{Hu}$,
Bosonic:	$\mathcal{O}_{H\square}, \mathcal{O}_{HG}, \mathcal{O}_{HW}, \mathcal{O}_{HB}, \mathcal{O}_W, \mathcal{O}_G,$
Yukawa:	$\mathcal{O}_{\tau H}, \mathcal{O}_{\mu H}, \mathcal{O}_{b H}, \mathcal{O}_{t H},$
Top 2F:	$\mathcal{O}_{HQ}^{(3)}, \mathcal{O}_{HQ}^{(1)}, \mathcal{O}_{Ht}, \mathcal{O}_{tG}, \mathcal{O}_{tW}, \mathcal{O}_{tB}, + \mathcal{O}_G$
Top 4F:	$\mathcal{O}_{Qq}^{3,1}, \mathcal{O}_{Qq}^{3,8}, \mathcal{O}_{Qq}^{1,8}, \mathcal{O}_{Qu}^8, \mathcal{O}_{Qd}^8, \mathcal{O}_{tQ}^8, \mathcal{O}_{tu}^8, \mathcal{O}_{td}^8.$


Individual limits: $U(3)^5$

2018 data: [Ellis et al.; JHEP 06 (2018) 146]




Zjj for triple gauge coupling

[ATLAS; CERN-EP-2020-045]





$\Delta\phi_{jj}$ distribution sensitive to linear C_W contributions


Wilson coefficient	Includes $ \mathcal{M}_{d6} ^2$	95% confidence interval [TeV $^{-2}$]	p -value (SM)
		Expected	Observed
c_W/Λ^2	no	[-0.30, 0.30]	45.9%
	yes	[-0.31, 0.29]	43.2%
\tilde{c}_W/Λ^2	no	[-0.12, 0.12]	82.0%
	yes	[-0.12, 0.12]	81.8%
c_{HWB}/Λ^2	no	[-2.45, 2.45]	29.0%
	yes	[-3.11, 2.10]	25.0%
$\tilde{c}_{HWB}/\Lambda^2$	no	[-1.06, 1.06]	1.7%
	yes	[-1.06, 1.06]	1.6%



Marginalised limits: U(3)⁵




Top-only: top + EWPO individual




- Some tension in $t\bar{t}$ data
- Asymmetries help to improve agreement

Top-only: breakdown



- $t\bar{t}$ asymmetries constrain orthogonal direction to cross section
- Large marginalisation effects: many similar operators
- $t\bar{t}V$ & $t\bar{t}H$ help to close the space
- Marginalised linear sensitivity: $C_{4F} \left[\frac{1 \text{ TeV}^2}{\Lambda^2} \right] \sim (5 - 15)$ significant $\frac{1}{\Lambda^4}$ effects

Top-only: top + EWPO marginalised




- C_{tH} : $t\bar{t}H$ bound alone is quite weak
- C_{tG} : Strong constraint but tension with SM
- Neutral top couplings poorly constrained
- EWPO closes $Zb\bar{b}$ coupling direction
- Impact of asymmetries in 4F
- Somewhat low scales (validity?)

Top-Higgs interplay

2D individual constraints






- All others set to 0
- $ggF/t\bar{t}H$ complementarity for (C_{HG}, C_{tH})
- H+jets STXS & $t\bar{t}V$ not yet competitive
- Strong impact of $t\bar{t}$ evident for (C_{tG}, C_G)
- Tension with SM $\sim 2\sigma$
- Significant correlations remain
- Large marginalisation effects (including 4F)

What is the concrete impact of 4F?



Individual 95% C. L.

- ggF+0 jet STXS
- $t\bar{t}H$
- ggF+ ≥ 1 jet STXS
- $t\bar{t}$
- $t\bar{t}V$
- Combined
- Marginalised



Point-like

Yukawa

Dipole

4F impact

Fit to ‘Higgs-only’ subspace

$$C_{H\square}, C_{HG}, C_{HW}, C_{HB}, C_{tH}, C_{bH}, C_{\tau H}, C_{\mu H} \\ + C_{tG} \& C_G$$

- Allow a closed fit to Higgs data only
- Emphasises impact of $t\bar{t}H$ & $t\bar{t}$

Now add in $t\bar{t}$ 4F operators


$$+ C_{Qq}^{3,8}, C_{Qq}^{1,8}, C_{Qu}^8, C_{Qd}^8, C_{tq}^8, C_{tu}^8, C_{td}^8$$

- Relatively mild impact
- Preferred $t\bar{t}$ phase space is different



C_{tG} : low $m_{t\bar{t}}$

4F : high $m_{t\bar{t}}$



- Able to constrain them independently



Full fit: individual



Full fit: marginalised



Correlations

Block diagonal:
correlations *within*
'sector'


Block off-diagonal:
correlations *among*
'sectors'

EWPO & top
~uncorrelated

EWPO-Higgs
 C_{HB} , C_{HW} , $C_{H\Box}$
& Yukawa
with EWPO


Higgs precision
rivaling LEP

Top-Higgs
 C_{HG} , C_G , C_{tH}
with 4F



PCA

2σ bound on Λ_i , $a_{ij}c_j = 1$




BSM implications

SMEFT-UV connection is model dependent by construction

- Implications on heavy new physics & validity of EFT is ***a posteriori***
- Depends on **sensitivity** & **energy scale** probed by data
- Bottom-up philosophy: new physics scale unknown

arbitrary dimensionful parameter $c_S = \frac{\lambda^2}{\Lambda^2}$ *coupling/mass scale of new physics*


constraint: $c/\Lambda^2 < X$



Difficult to address in a general way

- Today we are probing TeV scale new physics
- Hierarchies in sensitivity EWPO > Higgs > top (EW)
- Moderate-to-strong coupling scenarios most safe
- Generic NP in loops looks challenging for the LHC

BSM implications



Individual/marginalised = optimistic/pessimistic

- Real models should lie somewhere in between
- Less underlying parameters - more correlations
- Need to ‘re-run’ the fits to infer on underlying model parameters

Single field extensions

	Name	Spin	SU(3)	SU(2)	U(1)	Param.		Name	Spin	SU(3)	SU(2)	U(1)	Param.
Scalars	S	0	1	1	0	(M_S, κ_S)	Δ_1	$\frac{1}{2}$	1	2	$-\frac{1}{2}$	$(M_{\Delta_1}, \lambda_{\Delta_1})$	VLL
	S_1	0	1	1	1	(M_{S_1}, y_{S_1})	Δ_3	$\frac{1}{2}$	1	2	$-\frac{1}{2}$	$(M_{\Delta_3}, \lambda_{\Delta_3})$	
	φ	0	1	2	$\frac{1}{2}$	$(M_\varphi, Z_6 \cos \beta)$	Σ	$\frac{1}{2}$	1	3	0	$(M_\Sigma, \lambda_\Sigma)$	
	Ξ	0	1	3	0	(M_Ξ, κ_Ξ)	Σ_1	$\frac{1}{2}$	1	3	-1	$(M_{\Sigma_1}, \lambda_{\Sigma_1})$	
	Ξ_1	0	1	3	1	$(M_{\Xi_1}, \kappa_{\Xi_1})$	U	$\frac{1}{2}$	3	1	$\frac{2}{3}$	(M_U, λ_U)	
Z'	B	1	1	1	0	(M_B, \hat{g}_H^B)	D	$\frac{1}{2}$	3	1	$-\frac{1}{3}$	(M_D, λ_D)	VLL
	B_1	1	1	1	1	(M_{B_1}, g_{B_1})	Q_1	$\frac{1}{2}$	3	2	$\frac{1}{6}$	(M_{Q_1}, λ_{Q_1})	
	W	1	1	3	0	(M_W, \hat{g}_H^W)	Q_5	$\frac{1}{2}$	3	2	$-\frac{5}{6}$	(M_{Q_5}, λ_{Q_5})	
	W_1	1	1	3	1	$(M_{W_1}, \hat{g}_{W_1}^\varphi)$	Q_7	$\frac{1}{2}$	3	2	$\frac{7}{6}$	(M_{Q_7}, λ_{Q_7})	
	N	$\frac{1}{2}$	1	1	0	(M_N, λ_N)	T_1	$\frac{1}{2}$	3	3	$-\frac{1}{3}$	(M_{T_1}, λ_{T_1})	
VLQ	E	$\frac{1}{2}$	1	1	-1	(M_E, λ_E)	T_2	$\frac{1}{2}$	3	3	$\frac{2}{3}$	(M_{T_2}, λ_{T_2})	VLQ
	T	$\frac{1}{2}$	3	1	$\frac{2}{3}$	(M_T, s_L^t)	TB	$\frac{1}{2}$	3	2	$\frac{1}{6}$	$(M_{TB}, s_L^{t,b})$	

Considered single field extensions of the SM

- Complete tree-level matching dictionary is known [de Blas et al.; JHEP 03 (2018) 109]
- Interpret in terms of simplified 1 & 2 parameter versions of the models


One parameter models

Model	C_{HD}	C_{ll}	C_{Hl}^3	C_{Hl}^1	C_{He}	$C_{H\square}$	$C_{\tau H}$	C_{tH}	C_{bH}
S						$-\frac{1}{2}$			
S_1		1							
Σ			$\frac{1}{16}$	$\frac{3}{16}$			$\frac{y_\tau}{4}$		
Σ_1			$-\frac{1}{16}$	$-\frac{3}{16}$			$\frac{y_\tau}{8}$		
N			$-\frac{1}{4}$	$\frac{1}{4}$					
E			$-\frac{1}{4}$	$-\frac{1}{4}$			$\frac{y_\tau}{2}$		
Δ_1					$\frac{1}{2}$		$\frac{y_\tau}{2}$		
Δ_3					$-\frac{1}{2}$		$\frac{y_\tau}{2}$		
B_1	1					$-\frac{1}{2}$	$-\frac{y_\tau}{2}$	$-\frac{y_t}{2}$	$-\frac{y_b}{2}$
Ξ	-2					$\frac{1}{2}$	y_τ	y_t	y_b
W_1	$-\frac{1}{4}$					$-\frac{1}{8}$	$-\frac{y_\tau}{8}$	$-\frac{y_t}{8}$	$-\frac{y_b}{8}$
φ							$-y_\tau$	$-y_t$	$-y_b$
$\{B, B_1\}$						$-\frac{3}{2}$	$-y_\tau$	$-y_t$	$-y_b$
$\{Q_1, Q_7\}$								y_t	

Model	C_{Hq}^3	C_{Hq}^1	$(C_{Hq}^3)_{33}$	$(C_{Hq}^1)_{33}$	C_{Hu}	C_{Hd}	C_{tH}	C_{bH}	
U	$-\frac{1}{4}$	$\frac{1}{4}$	$-\frac{1}{4}$	$\frac{1}{4}$			$\frac{y_t}{2}$		
D	$-\frac{1}{4}$	$-\frac{1}{4}$	$-\frac{1}{4}$	$-\frac{1}{4}$				$\frac{y_b}{2}$	
Q_5						$-\frac{1}{2}$		$\frac{y_b}{2}$	
Q_7					$\frac{1}{2}$		$\frac{y_t}{2}$		
T_1	$-\frac{1}{16}$	$-\frac{3}{16}$	$-\frac{1}{16}$	$-\frac{3}{16}$			$\frac{y_t}{4}$	$\frac{y_b}{8}$	
T_2	$-\frac{1}{16}$	$\frac{3}{16}$	$-\frac{1}{16}$	$\frac{3}{16}$			$\frac{y_t}{8}$	$\frac{y_b}{4}$	
T			$-\frac{1}{2} \frac{M_T^2}{v^2}$	$\frac{1}{2} \frac{M_T^2}{v^2}$			$y_t \frac{M_T^2}{v^2}$		

$$\times \frac{\lambda^2}{M^2}$$

One parameter models



Tree-level patterns

Similar particles often generate similar operator patterns

- e.g. Massive vector bosons B_1, W_1 : $C_{H\square} = C_{tH} = \pm \frac{1}{2} C_{HD}$
- Study pattern-inspired subspaces


Boson-specific: $(C_{HD}, C_{H\square}, C_{tH})$,

Lepton-specific: $(C_{He}, C_{H\ell}^{(1,3)}, C_{\ell\ell})$,


Quark-specific: $(C_{Hu}, C_{Hd}, C_{Hq}^{(1,3)}, C_{tH})$,

Top-specific: $((C_{Hq}^{(1)})_{33}, (C_{Hq}^{(3)})_{33}, C_{HG}, C_{bH}, C_{tH}, C_{Ht})$

Boson specific: $C_{H\square}, C_{HD}, C_{tH}$



Boson specific




Pull-ology

Brute force: fit to all combinations of n-coefficients

$$P \equiv \sqrt{\chi^2_{SM} - \chi^2_{\text{fit}}}$$

- Agnostic search for improved fit w.r.t SM
- NP hints could show up in this way
- Advantage of fast, linear fit method
- Highlights tension in $t\bar{t}$ data
- No conclusive NP hints so far...



Conclusions

Presented first EWPO, Higgs, Diboson & Top fit in SMEFT

- Include leading contributions from top operators in ggF
- Top & Higgs sector are starting to talk to each other
- $t\bar{t}$ 4 fermion operators don't appear to spoil naive picture of interplay

Analytic, linear analysis has many benefits

- Simple likelihood described by best fit+correlations, PCA exact
- Easy to interpret/combine with other likelihoods
- Fast: repeat for subsets, BSM interpretations

& Drawbacks

- Potentially important quadratic effects, especially in top data
- Gaussian priors only, not really appropriate for th. errors

Outlook

Much more to be done

- Explore the likelihood further
- Compare results to a quadratic fit to test validity
- SMEFT theory errors?
- Explore impact of new observables: VBS, VVV, rare top modes



Impact of loops

- Top operators in loops: Higgs decays + EWPO
- NLO corrections


BSM implications

- Go beyond 1 particle benchmarks towards realistic models
- Are there compelling top/Higgs scenarios that admit a valid EFT interpretation with LHC data?


Bonus: di-Higgs in SMEFT



$$\begin{aligned}
 O_\phi &= (H^\dagger H)^3 \\
 O_{tG} &= (\bar{Q}\sigma_{\mu\nu}T^A t) \tilde{H} G_A^{\mu\nu} \\
 O_{t\phi} &= (H^\dagger H)(\bar{Q}t \tilde{H}) \\
 O_{d\phi} &= \partial_\mu(H^\dagger H)\partial^\mu(H^\dagger H) \\
 O_{\phi G} &= (H^\dagger H)G_A^{\mu\nu}G_{\mu\nu}^A
 \end{aligned}$$



Backup



Data: EWPO & Diboson

EW precision observables	n_{obs}
Precision electroweak measurements on the Z resonance.	12
Γ_Z , $\sigma_{\text{had.}}^0$, R_ℓ^0 , A_{FB}^ℓ , $A_\ell(\text{SLD})$, $A_\ell(\text{Pt})$, R_b^0 , R_c^0 , A_{FB}^b , A_{FB}^c , A_b & A_c	
Combination of CDF and D0 W -Boson Mass Measurements	1
LHC run 1 W boson mass measurement by ATLAS	1
Diboson LEP & LHC	n_{obs}
$W^+ W^-$ angular distribution measurements at LEP II.	8
$W^+ W^-$ total cross section measurements at L3 in the $\ell\nu\ell\nu$, $\ell\nu qq$ & $qqqq$ final states for 8 energies	24
$W^+ W^-$ total cross section measurements at OPAL in the $\ell\nu\ell\nu$, $\ell\nu qq$ & $qqqq$ final states for 7 energies	21
$W^+ W^-$ total cross section measurements at ALEPH in the $\ell\nu\ell\nu$, $\ell\nu qq$ & $qqqq$ final states for 8 energies	21
ATLAS $W^+ W^-$ differential cross section in the $e\nu\mu\nu$ channel, $\frac{d\sigma}{dp_{\ell_1}^T}$, $p_T > 120$ GeV overflow bin	1
ATLAS $W^+ W^-$ fiducial differential cross section in the $e\nu\mu\nu$ channel, $\frac{d\sigma}{dp_{\ell_1}^T}$	14
ATLAS $W^\pm Z$ fiducial differential cross section in the $\ell^+\ell^-\ell^\pm\nu$ channel, $\frac{d\sigma}{dp_Z^T}$	7
CMS $W^\pm Z$ normalised fiducial differential cross section in the $\ell^+\ell^-\ell^\pm\nu$ channel, $\frac{1}{\sigma} \frac{d\sigma}{dp_Z^T}$	11
ATLAS Zjj fiducial differential cross section in the $\ell^+\ell^-$ channel, $\frac{d\sigma}{d\Delta\varphi_{jj}}$	12

Data: Higgs

LHC Run 1 Higgs	n_{obs}
ATLAS and CMS LHC Run 1 combination of Higgs signal strengths. Production: ggF , VBF , ZH , WH & tH Decay: $\gamma\gamma$, ZZ , W^+W^- , $\tau^+\tau^-$ & $b\bar{b}$	21
ATLAS inclusive $Z\gamma$ signal strength measurement	1
LHC Run 2 Higgs (new)	n_{obs}
ATLAS combination of signal strengths and stage 1.0 STXS in $H \rightarrow 4\ell$ including ratios of branching fractions to $\gamma\gamma$, WW^* , $\tau^+\tau^-$ & $b\bar{b}$ Signal strengths coarse STXS bins fine STXS bins	16 19 25
CMS LHC combination of Higgs signal strengths. Production: ggF , VBF , ZH , WH & tH Decay: $\gamma\gamma$, ZZ , W^+W^- , $\tau^+\tau^-$, $b\bar{b}$ & $\mu^+\mu^-$	23
CMS stage 1.0 STXS measurements for $H \rightarrow \gamma\gamma$. 13 parameter fit 7 parameter fit	13 7
CMS stage 1.0 STXS measurements for $H \rightarrow \tau^+\tau^-$	9
CMS stage 1.1 STXS measurements for $H \rightarrow 4\ell$	19
CMS differential cross section measurements of inclusive Higgs production in the $WW^* \rightarrow \ell\nu\ell\nu$ final state.	5 6
$\frac{d\sigma}{dn_{\text{jet}}}$ $\frac{d\sigma}{dp_H^T}$	
ATLAS $H \rightarrow Z\gamma$ signal strength.	1
ATLAS $H \rightarrow \mu^+\mu^-$ signal strength.	1

Data: Tevatron, LHC Run 1 & 2 top



Tevatron & Run 1 top	n_{obs}	Ref.	Run 2 top	n_{obs}	Ref.
Tevatron combination of differential $t\bar{t}$ forward-backward asymmetry, $A_{FB}(m_{t\bar{t}})$.	4	[7]	CMS $t\bar{t}$ differential distributions in the dilepton channel. $\frac{d\sigma}{dm_{t\bar{t}}}$	6	[46, 50]
ATLAS $t\bar{t}$ differential distributions in the dilepton channel. $\frac{d\sigma}{dm_{t\bar{t}}}$	6	[31]	CMS $t\bar{t}$ differential distributions in the $\ell+\text{jets}$ channel. $\frac{d\sigma}{dm_{t\bar{t}}}$	10	[53]
ATLAS $t\bar{t}$ differential distributions in the $\ell+\text{jets}$ channel. $\frac{d\sigma}{dm_{t\bar{t}}} \mid \frac{d\sigma}{d y_{t\bar{t}} } \mid \frac{d\sigma}{dp_t^T} \mid \frac{d\sigma}{d y_t }$.	7 5 8 5	[24]	ATLAS measurement of differential $t\bar{t}$ charge asymmetry, $A_C(m_{t\bar{t}})$. $\frac{d\sigma}{dp_z^T} \mid \frac{d\sigma}{d\cos\theta^*}$	5	[55]
CMS $t\bar{t}$ differential distributions in the $\ell+\text{jets}$ channel. $\frac{d\sigma}{dm_{t\bar{t}}} \mid \frac{d\sigma}{d y_{t\bar{t}} } \mid \frac{d\sigma}{dp_t^T} \mid \frac{d\sigma}{d y_t }$.	7 10 8 10	[25, 34]	ATLAS $t\bar{t}W$ & $t\bar{t}Z$ cross section measurements. $\sigma_{t\bar{t}W} \mid \sigma_{t\bar{t}Z}$	2	[58]
CMS measurement of differential $t\bar{t}$ charge asymmetry, $A_C(m_{t\bar{t}})$ in the dilepton channel.	3	[33]	CMS $t\bar{t}W$ & $t\bar{t}Z$ cross section measurements. $\sigma_{t\bar{t}W} \mid \sigma_{t\bar{t}Z}$	1 1	[48]
ATLAS inclusive measurement $t\bar{t}$ charge asymmetry, $A_C(m_{t\bar{t}})$ in the dilepton channel.	1	[32]	CMS $t\bar{t}Z$ differential distributions. $\frac{d\sigma}{dp_\gamma^T}$	4 4	[60]
ATLAS & CMS combination of differential $t\bar{t}$ charge asymmetry, $A_C(m_{t\bar{t}})$, in the $\ell+\text{jets}$ channel.	6	[38]	ATLAS $t\bar{t}\gamma$ differential distribution. $\frac{d\sigma}{dp_{t+\bar{t}}^T} \mid R_t(p_{t+\bar{t}}^T)$	11	[62]
CMS $t\bar{t}$ double differential distributions in the dilepton channel. $\frac{d\sigma}{dm_{t\bar{t}} dy_t} \mid \frac{d\sigma}{dm_{t\bar{t}} d y_{t\bar{t}} } \mid \frac{d\sigma}{dm_{t\bar{t}} dp_{t\bar{t}}^T} \mid \frac{d\sigma}{dy_t dp_t^T}$.	16 16 16 16	[18, 35]	CMS measurement of differential cross sections and charge ratios for t -channel single-top quark production. $\sigma_t \mid \sigma_{\bar{t}} \mid \sigma_{t+\bar{t}} \mid R_t$.	5 5	[56]
ATLAS & CMS Run 1 combination of W -boson helicity fractions in top decay. f_0, f_L & f_R	3	[40]	CMS measurement of t -channel single-top and anti-top cross sections. $\sigma_t, \sigma_{\bar{t}}, \sigma_{t+\bar{t}} \& R_t$.	4	[42]
ATLAS measurement of W -boson helicity fractions in top decay. f_0, f_L & f_R	3	[30]	CMS measurement of the t -channel single-top and anti-top cross sections. $\sigma_t \mid \sigma_{\bar{t}} \mid \sigma_{t+\bar{t}} \mid R_t$.	1 1 1 1	[45]
CMS measurement of W -boson helicity fractions in top decay. f_0, f_L & f_R	3	[29]	CMS t -channel single-top differential distributions. $\frac{d\sigma}{dp_{t+\bar{t}}^T} \mid \frac{d\sigma}{d y_{t+\bar{t}} }$	4 4	[44]
ATLAS $t\bar{t}W$ & $t\bar{t}Z$ cross section measurements. $\sigma_{t\bar{t}W} \mid \sigma_{t\bar{t}Z}$	2	[23]	ATLAS tW cross section measurement.	1	[43]
CMS $t\bar{t}W$ & $t\bar{t}Z$ cross section measurements. $\sigma_{t\bar{t}W} \mid \sigma_{t\bar{t}Z}$	2	[26]	CMS tZ cross section measurement.	1	[47]
ATLAS $t\bar{t}\gamma$ cross section measurement in the $\ell+\text{jets}$ channel.	1	[36]	CMS tW cross section measurement.	1	[52]
CMS $t\bar{t}\gamma$ cross section measurement in the $\ell+\text{jets}$ channel.	1	[37]	ATLAS tZ cross section measurement.	1	[49]
ATLAS t -channel single-top differential distributions. $\frac{d\sigma}{dp_t^T} \mid \frac{d\sigma}{dp_{\bar{t}}^T} \mid \frac{d\sigma}{d y_t } \mid \frac{d\sigma}{d y_{\bar{t}} }$	4 4 4 5	[39]	CMS tZ ($Z \rightarrow \ell^+ \ell^-$) cross section measurement	1	[54]
CMS s -channel single-top cross section measurement.	1	[28]	ATLAS four-top search in the multi-lepton and same-sign dilepton channels.	1	[63]
CMS t -channel single-top differential distributions. $\frac{d\sigma}{dp_{t+\bar{t}}^T} \mid \frac{d\sigma}{d y_{t+\bar{t}} }$	6 6	[19]	ATLAS four-top search in the single-lepton and opposite-sign dilepton channels.	1	[51]
CMS measurement of the t -channel single-top and anti-top cross sections. $\sigma_t \mid \sigma_{\bar{t}} \mid \sigma_{t+\bar{t}} \mid R_t$.	1 1 1 1	[20]	CMS four-top search in the multi-lepton and same-sign dilepton channels.	1	[61]
ATLAS s -channel single-top cross section measurement.	1	[27]	CMS four-top search in the single-lepton and opposite-sign dilepton channels.	1	[59]
CMS tW cross section measurement.	1	[21]	CMS $t\bar{t}b\bar{b}$ cross section measurement in the all-jet channel.	1	[57]
ATLAS tW cross section measurement in the single lepton channel.	1	[41]	CMS $t\bar{t}b\bar{b}$ cross section measurement in the dilepton channel.	1	[64]
ATLAS tW cross section measurement in the dilepton channel.	1	[22]			

Fisher information breakdown



C_i	EWPO	LEP WW	Run 1 SS	Run 2 SS	STXS	LHC WW	WZ	Zjj	$t\bar{t}$	$W_{\text{hel.}}$	tX	$t\bar{t}V$
C_{HWB}	51	—	7	14	28	—	—	—	—	—	—	—
C_{HD}	100	—	—	—	—	—	—	—	—	—	—	—
C_{ll}	99	—	—	—	—	—	—	—	—	—	—	—
$C_{Hl}^{(3)}$	99	—	—	—	—	—	—	—	—	—	—	—
$C_{Hl}^{(1)}$	100	—	—	—	—	—	—	—	—	—	—	—
C_{He}	100	—	—	—	—	—	—	—	—	—	—	—
$C_{Hq}^{(3)}$	89	1	—	—	2	—	6	—	—	—	—	—
$C_{Hq}^{(1)}$	99	—	—	—	—	—	—	—	—	—	—	—
C_{Hd}	99	—	—	—	—	—	—	—	—	—	—	—
C_{Hu}	98	—	—	—	1	—	—	—	—	—	—	—
$C_{H\square}$	—	—	22	46	32	—	—	—	—	—	—	—
C_{HG}	—	—	22	42	36	—	—	—	—	—	—	—
C_{HW}	—	—	14	29	56	—	—	—	—	—	—	—
C_{HB}	—	—	14	29	57	—	—	—	—	—	—	—
C_W	—	3	—	—	—	—	13	84	—	—	—	—
C_G	—	—	—	—	—	—	—	—	43	—	—	56
$C_{\tau H}$	—	—	22	45	34	—	—	—	—	—	—	—
$C_{\mu H}$	—	—	5	95	—	—	—	—	—	—	—	—
C_{bH}	—	—	19	35	47	—	—	—	—	—	—	—
C_{tH}	—	—	21	45	34	—	—	—	—	—	—	—
$C_{HQ}^{(3)}$	99	—	—	—	—	—	—	—	—	—	—	—
$C_{HQ}^{(1)}$	100	—	—	—	—	—	—	—	—	—	—	—
C_{Ht}	—	—	—	—	—	—	—	—	—	—	—	100
C_{tG}	—	—	13	29	24	—	—	—	24	—	—	9
C_{tW}	—	—	—	—	—	—	—	—	—	84	15	—
C_{tB}	—	—	—	—	—	—	—	—	—	—	—	100
$C_{Qq}^{3,1}$	—	—	—	—	—	—	—	—	—	—	100	—
$C_{Qq}^{3,8}$	—	—	—	—	—	—	—	—	87	—	—	13
$C_{Qq}^{1,8}$	—	—	—	—	—	—	—	—	82	—	—	17
C_{Qu}^8	—	—	—	—	—	—	—	—	91	—	—	7
C_{Qd}^8	—	—	—	2	—	—	—	—	92	—	—	6
C_{tq}^8	—	—	—	1	—	—	—	—	89	—	—	10
C_{tu}^8	—	—	—	—	—	—	—	—	96	—	—	3
C_{td}^8	—	—	—	2	—	—	—	—	92	—	—	5

Removing C_G

With



Without



Removing C_G

